

PAM Health

THE ROAD TO RECOVERY BEGINS HERE



PAM Health

60% CMS-13 Diagnoses

Medicare requires that 60% of all patients admitted to an Acute Rehabilitation Hospital must have a primary diagnosis that falls within 1 of 13 different CMS-13 categories.

STROKE/CVA

- Stroke/CVA or late effect CVA (w/hemiparesis)
- Cerebral embolism/thrombosis
- Intracerebral bleed
- Hemiparesis related to stroke/CVA
- Hypertensive encephalopathy
- Moyamoya disease

SPINAL CORD INJURY (Non-Traumatic and Traumatic)

- Malignant neoplasm of the spine
- Neoplasm of the spinal cord/meninges
- Encephalitis/myelitis/encephalomyelitis
- Para/Quadriplegia due to spinal cord injury
- Caudal equina syndrome
- Intraspinal abscess
- Specified myelopathies or spinal stroke
- Paraplegia, quadriplegia
- Fracture of vertebrae with spinal cord injury
- Late effect spinal cord injury
- Cervical or lumbar disc disorder with myelopathy



THE ROAD TO RECOVERY BEGINS HERE

CONGENITAL DEFORMITY

- Dwarfism
- Acquired torsion dystonia
- Cerebral palsy
- Spinal muscular atrophy
- Congenital deformity of the hip
- Congenital reduction of leg/arm
- Congenital absence of leg/arm
- Congenital scoliosis

AMPUTATIONS

- Traumatic amputation of leg or foot
- Complication of amputated stump
- Acquired amputation of leg or foot
- Prosthetic training s/p recent amputation

MAJOR MULTIPLE TRAUMA

- Brain + spinal cord injury
- Brain + multiple fxs/amputation
- Spinal cord + multiple fxs/amputation
- Fx of both legs
- Fx of leg + arm and/or a rib sternum
- Disruption of the pelvic circle

FRACTURE OF HIP

- Pathological fx of the hip/femoral neck
- Traumatic fx of the hip/femoral neck or acetabulum
- Trochanteric fx of the femur

BRAIN INJURY (Non-Traumatic and Traumatic)

- Traumatic bleed of the brain or late effects
- Post-concussion syndrome
- Meningitis (currently on antibiotics)
- Specified encephalitis or late effects
- Specified encephalopathy
- Neoplasms of the brain or pituitary gland
- Subarachnoid and subdural hemorrhage
- Fracture of the skull vault
- Concussion with loss of consciousness
- Cerebral contusion or late effects

NEUROLOGICAL DISORDERS

- Parkinson's disease
- Specified types of polyneuropathy
- Post-polio syndrome
- Spinal muscular dystrophy
- Multiple sclerosis
- Neuromyelitis
- Brachial, cervical, lumbosacral or thoracic root nerve or plexus disorders
- Guillain-Barré
- Myasthenia gravis
- Specified myopathies and polymyositis

BURNS

- Late effects: cicatrix, deformity, and/or scarring that prevents normal function of extremities

SYSTEMIC VASCULITIDIES

- Systemic lupus erythematosus

KNEE OR HIP REPLACEMENT

- Bilateral knees or hips; BMI>50; age 85 or older

PATIENT CONSIDERATIONS

- Prior level of independence
- Functional impairments affecting mobility, activities of daily living, swallowing or cognition
- Late effects from a previous neurological injury or illness, i.e. brain injury, CVA or SCI
- Co-morbid conditions or complications
- Supporting physician, nursing, and therapy documentation

40% OTHER REHAB DIAGNOSES

Neurological Disorders

- Chemo or radiation

Induced Neuropathy

- Neuropathy
- Polyneuropathy

Orthopedic Injuries

- Contusions
- Non-operative fractures
- Pelvic fractures
- Severe joint injuries
- Upper extremity fractures

Back Diagnosis

- Compression fractures
- Degenerative disc disease
- Degenerative joint disease
- Laminectomy
- Radiculopathy
- Spinal stenosis

Cardiac Conditions

- Abdominal aortic aneurysm
- Coronary artery bypass graft
- Congestive heart failure
- Cardiac arrhythmia
- Myocardial infarction

Respiratory (non-ventilator)

- COPD exacerbation
- Pneumonia
- Respiratory infections
- Respiratory failure

Deconditioning Diagnosis

- Decreased endurance and strength
- Debility
- Frequent falls
- Unsafe transfers
- Unsteady gait

Other

- Chronic pain
- Cellulitis
- Peripheral vascular disease
- Urinary tract infections
- Wound care

